

Treatment Options by Symptom Severity¹

EPDS Score or Clinical **Assessment**

EPDS 0-8

EPDS 9-13

EPDS 14-18

EPDS ≥19

Treatment Options

Treatment options in each column may overlap

Limited To No Symptoms	Mild Symptoms	Moderate Symptoms	Severe Symptoms
		Consider inpatient hospitalization when safety or ability to care for self or baby is a concern	Strongly consider inpatient hospitalization when safety or ability to care for self or baby is a concern
	May consider medication	Strongly consider medication	Strongly consider medication
Therapy for mother	Therapy for mother	Therapy for mother	Therapy for mother
Dyadic therapy for mother/baby			
Community/social support (including support groups)			
Consider as augmentation: Complementary/ Alternative therapies (bright light therapy, Omega-3 fatty acids, acupuncture, folate, massage)	Consider as augmentation: Complementary/ Alternative therapies (bright light therapy, Omega-3 fatty acids, acupuncture, folate, massage)	Consider as augmentation: Complementary/ Alternative therapies (bright light therapy, Omega-3 fatty acids, acupuncture, folate, massage)	Consider as augmentation: Complementary/ Alternative therapies (bright light therapy, Omega-3 fatty acids, acupuncture, folate, massage)
Support with dysregulated baby; crying, sleep, feeding problems	Support with dysregulated baby; crying, sleep, feeding problems	Support with dysregulated baby; crying, sleep, feeding problems	Support with dysregulated baby; crying, sleep, feeding problems
Physical activity	Physical activity	Physical activity	Physical activity
Self-care (sleep, hygiene, healthy diet) and lactation support, if needed	Self-care (sleep, hygiene, healthy diet)	Self-care (sleep, hygiene, healthy diet)	Self-care (sleep, hygiene, healthy diet)

Information adapted from: Montgomery SA, Asberg M: A new depression scale designed to be sensitive to change. British Journal of Psychiatry 134:382-389, 1979

This resource has been adapted with permission from the MCPAP for Moms Provider Toolkit.